

On graded irreducible representations of Leavitt path algebras

Roozbeh Hazrat^(a) and Kulumani M. Rangaswamy^(b)

(a) Centre for Research in Mathematics,
University of Western Sydney, Sydney, Australia
E-mail: r.hazrat@uws.edu.au

(b) Department of Mathematics,
University of Colorado at Colorado Springs,
Colorado Springs, Colorado 80918, USA
E-mail: krangasw@uccs.edu

Abstract

Using the E -algebraic branching systems, various graded irreducible representations of a Leavitt path K -algebra L of a directed graph E are constructed. The concept of a Laurent vertex is introduced and it is shown that the minimal graded left ideals of L are generated by the Laurent vertices or the line points leading to a detailed description of the graded socle of L . Following this, a complete characterization is obtained of the Leavitt path algebras over which every graded irreducible representation is finitely presented. A useful result is that the irreducible representation $V_{[p]}$ induced by infinite paths tail-equivalent to an infinite path p (we call this a Chen simple module) is graded if and only if p is an irrational path. We also show that every one-sided ideal of L is graded if and only if the graph E contains no cycles. Supplementing the theorem of one of the co-authors that every Leavitt path algebra L is graded von Neumann regular, we show that L is graded self-injective if and only if L is a graded semi-simple algebra, made up of matrix rings of arbitrary size over the field K or the graded field $K[x^n, x^{-n}]$ where $n \in \mathbb{N}$.

1 Introduction and Preliminaries

As mentioned in [16], branching systems arise in the study of various disciplines such as random walk, symbolic dynamics and scientific computing (see e.g. [15], [22], [28]). Given an arbitrary directed graph E , Gonçalves and Royer defined in [16] and [17] a branching system using a measure space (X, μ) and indicated a method of constructing a large number of representations of the graph C^* -algebra $C^*(E)$ in the space of bounded linear operators on $L^2(X, \mu)$. As an algebraic analogue of this theory, they defined the concept of an E -algebraic

branching system in [18] to construct various representations of Leavitt path algebras (which are algebraic analogues of graph C^* -algebras). Following this, X.W.Chen ([14]) constructed irreducible representations of a Leavitt path algebra L of a graph E using the sinks as well as infinite paths in E and showed that these representations are also induced by appropriate E -algebraic branching systems. Just about the same time, the relations between the theory of quiver representations and the theory of representations of Leavitt path algebras were explored in [9]. Following this there was a flurry of activity in investigating various irreducible representations of L induced by infinite emitters and cycles in E , leading to the study of Leavitt path algebras with a special type of or a specific number of irreducible representations (see [6], [11], [26], [27]).

Leavitt path algebras are also endowed with a compatible $\mathbb{Z}$ -grading and the initial study of the graded structure of these algebras was done in [20]. Moreover, it was shown in [21] that every Leavitt path algebra is a graded von Neumann regular algebra, an indication of a possible rich internal structure of these algebras. As a first step in understanding the graded modules over these algebras, we consider in this paper the graded irreducible representations of a Leavitt path algebra L of an arbitrary graph E . It is known for a graded von Neumann regular ring R with identity, that there is a one-to-one correspondence between the graded minimal left ideals of R and the minimal left ideals of R_0 . When R is a Leavitt path algebra, R_0 is ultramatricial. While it may not always be possible to describe all the minimal left ideals of R_0 , our graphical approach enables us to provide a complete description of the graded minimal left/right ideals of R . Defining a grade on the E -algebraic branching systems, we construct various types of graded irreducible representations of L . A useful result is that if p is an infinite path, then the irreducible representation $V_{[p]}$ defined by the equivalence class of infinite paths tail-equivalent to p is graded if and only if p is an irrational path (see definitions below). We also describe the annihilator ideals of many of the graded irreducible representations. It turns out that, unlike the non-graded case, the annihilator of a graded irreducible representation need not be a primitive ideal, but is a graded prime ideal of L . The concept of a Laurent vertex (see Definition 2.6 below) is introduced and is shown that a graded left ideal of L is a minimal graded left ideal if and only if it is isomorphic to the left ideal Lv with possibly a shifted grading, where v is either a Laurent vertex or a line point. This enables us to describe the graded socle of L as the ideal generated by these special vertices in E and that it is a graded direct sum of matrix rings of arbitrary size over K and/or $K[x, x^{-1}]$ with appropriate gradings.

Next we investigate when a graded irreducible representation of L is finitely presented. A key result is that a graded irreducible representation S induced by an infinite path p is finitely presented if and only if p contains a line point and in that case S becomes graded projective. Using this, it is shown that every irreducible representation of L is finitely presented if and only if the graph E is row-finite and that every path in E eventually ends at a line point or a Laurent vertex. In this case, the structure of L is also completely described.

A well-known theorem in Leavitt path algebras ([29]) states that every two-

sided ideal of a Leavitt path algebra $L := L_K(E)$ is graded if and only if the graph E satisfies Condition (K) (see definition below). A natural question is when will every left (right) ideal of L be graded. We show that this happens exactly when the graph E contains no cycles. In this case, L is a directed union of graded subalgebras B each of which is a graded direct sum of finitely many matrix rings of finite order over K endowed with natural gradings.

It is known that Leavitt path algebras which are self-injective are always von Neumann regular (see [13]). Since by [21] every Leavitt path algebra is graded von Neumann regular, it is natural to consider the subclass of Leavitt path algebras which are graded self-injective. We show that such algebras are just the graded semi-simple K -algebras. The corresponding graph theoretical properties of the graph E are also described.

As an illustration of the advantage of viewing a Leavitt path algebra L as a $\mathbb{Z}$ -graded algebra, we show how the properties of L as a graded structure enables us to simplify, generalize or correct the proofs of some of the important previously published results on Leavitt path algebras (see Theorems 4.2, 4.6 and Proposition 6.2).

For the general notation, terminology and results in Leavitt path algebras, we refer to [1] and [25]. We give below a short outline of some of the needed basic concepts and results.

A (directed) graph $E = (E^0, E^1, r, s)$ consists of two sets E^0 and E^1 together with maps $r, s : E^1 \rightarrow E^0$. The elements of E^0 are called *vertices* and the elements of E^1 *edges*. All the graphs E that we consider (excepting when specifically stated) are arbitrary in the sense that no restriction is placed either on the number of vertices in E or on the number of edges emitted by a single vertex. In this connection, we wish to point out that some of the earlier papers in Leavitt path algebras, such as [2], [12] which we quote in this paper, assume that the graphs considered are countable. But the proofs of the results in these papers are valid even for uncountable graphs.

A vertex v is called a *sink* if it emits no edges and a vertex v is called a *regular vertex* if it emits a non-empty finite set of edges. An *infinite emitter* is a vertex which emits infinitely many edges. A graph without infinite emitters is said to be row-finite. For each $e \in E^1$, we call e^* a ghost edge. We let $r(e^*)$ denote $s(e)$, and we let $s(e^*)$ denote $r(e)$. A *path* μ of length $n > 0$ is a finite sequence of edges $\mu = e_1 e_2 \cdots e_n$ with $r(e_i) = s(e_{i+1})$ for all $i = 1, \dots, n-1$. In this case $\mu^* = e_n^* \cdots e_2^* e_1^*$ is the corresponding ghost path. A vertex is considered a path of length 0. The set of all vertices on the path μ is denoted by μ^0 . We shall denote the length of a path μ by $|\mu|$.

A path $\mu = e \cdots e_n$ in E is *closed* if $r(e_n) = s(e_1)$, in which case μ is said to be based at the vertex $s(e_1)$. A closed path μ as above is called *simple* provided it does not pass through its base more than once, i.e., $s(e_i) \neq s(e_1)$ for all $i = 2, \dots, n$. The closed path μ is called a *cycle* if it does not pass through any of its vertices twice, that is, if $s(e_i) \neq s(e_j)$ for every $i \neq j$. An *exit* for a path $\mu = e_1 \cdots e_n$ is an edge f that satisfies $s(f) = s(e_i)$ for some i but $f \neq e_i$. The graph E is said to satisfy *Condition (L)*, if every cycle in E has an exit in

E . The graph E is said to satisfy *Condition (K)*, if any vertex on a closed path μ is also the base for a closed path γ different from μ .

If there is a path from vertex u to a vertex v , we write $u \geq v$. A subset D of vertices is said to be *downward directed* if for any $u, v \in D$, there exists a $w \in D$ such that $u \geq w$ and $v \geq w$. A subset H of E^0 is called *hereditary* if, whenever $v \in H$ and $w \in E^0$ satisfy $v \geq w$, then $w \in H$. A hereditary set is *saturated* if, for any regular vertex v , $r(s^{-1}(v)) \subseteq H$ implies $v \in H$.

Given an arbitrary graph E and a field K , the *Leavitt path algebra* $L_K(E)$ is defined to be the K -algebra generated by a set $\{v : v \in E^0\}$ of pairwise orthogonal idempotents together with a set of variables $\{e, e^* : e \in E^1\}$ which satisfy the following conditions:

- (1) $s(e)e = e = er(e)$ for all $e \in E^1$.
- (2) $r(e)e^* = e^* = e^*s(e)$ for all $e \in E^1$.
- (3) (The "CK-1 relations") For all $e, f \in E^1$, $e^*e = r(e)$ and $e^*f = 0$ if $e \neq f$.
- (4) (The "CK-2 relations") For every regular vertex $v \in E^0$,

$$v = \sum_{e \in E^1, s(e)=v} ee^*.$$

For any vertex v , the *tree* of v is $T_E(v) = \{w \in E^0 : v \geq w\}$. We say there is a *bifurcation* at a vertex v or v is a *bifurcation vertex*, if v emits more than one edge. In a graph E , a vertex v is called a *line point* if there is no bifurcation or a cycle based at any vertex in $T_E(v)$. Thus, if v is a line point, there will be a single finite or infinite line segment μ starting at v (μ could just be v) and any other path α with $s(\alpha) = v$ will just be an initial subpath of μ . It was shown in [12] that v is a line point in E if and only if $L_K(E)v$ (and likewise $vL_K(E)$) is a simple left (right) ideal of L .

We shall be using the following concepts and results from [29]. A *breaking vertex* of a hereditary saturated subset H is an infinite emitter $w \in E^0 \setminus H$ with the property that $0 < |s^{-1}(w) \cap r^{-1}(E^0 \setminus H)| < \infty$. The set of all breaking vertices of H is denoted by B_H . For any $v \in B_H$, v^H denotes the element $v - \sum_{s(e)=v, r(e) \notin H} ee^*$. Given a hereditary saturated subset H and a subset $S \subseteq B_H$, (H, S) is called an *admissible pair*. Given an admissible pair (H, S) , the ideal generated by $H \cup \{v^H : v \in S\}$ is denoted by $I(H, S)$. It was shown in [29] that the graded ideals of $L_K(E)$ are precisely the ideals of the form $I(H, S)$ for some admissible pair (H, S) . Moreover, $L_K(E)/I(H, S) \cong L_K(E \setminus (H, S))$. Here $E \setminus (H, S)$ is the *quotient graph* of E in which

$$(E \setminus (H, S))^0 = (E^0 \setminus H) \cup \{v' : v \in B_H \setminus S\}$$

and

$$(E \setminus (H, S))^1 = \{e \in E^1 : r(e) \notin H\} \cup \{e' : e \in E^1, r(e) \in B_H \setminus S\}.$$

Further, r and s are extended to $(E \setminus (H, S))^1$ by setting $s(e') = s(e)$ and $r(e') = r(e)'$.

A useful observation is that every element a of $L_K(E)$ can be written as $a = \sum_{i=1}^n k_i \alpha_i \beta_i^*$, where $k_i \in K$, α_i, β_i are paths in E and n is a suitable integer.

Moreover, $L_K(E) = \bigoplus_{v \in E^0} L_K(E)v = \bigoplus_{v \in E^0} vL_K(E)$ (see [1]).

Recall that a K -algebra R is said to be a $\mathbb{Z}$ -graded algebra if $R = \bigoplus_{n \in \mathbb{Z}} R_n$ is a direct sum of K -vector spaces R_n , which satisfy the property that $R_m R_n \subseteq R_{m+n}$ for all m, n . Elements of the subspaces R_n are called *homogeneous elements*. A left R -module M is called a *graded left R -module*, if $M = \bigoplus_{n \in \mathbb{Z}} M_n$ is

a direct sum of subgroups M_n which satisfy the property that $R_m M_n \subseteq M_{m+n}$ for all m, n . For any $k \in \mathbb{Z}$, the *k -shifted module* $M(k)$ is the graded R -module for which the homogeneous component $M(k)_n = M_{k+n}$ for all n . A graded ring R with 1 is called a *graded division ring* if every non-zero homogeneous element in R is invertible. A commutative graded division ring is called a *graded field*. The graded fields that we encounter in this paper are $\mathbb{Z}$ -graded subrings of $K[x, x^{-1}]$ of the form $K[x^n, x^{-n}]$ with support $n\mathbb{Z}$, where $n \in \mathbb{N}$. For the general properties of graded rings and graded modules, we refer to [24]. Every Leavitt path algebra $L := L_K(E)$ is a $\mathbb{Z}$ -graded algebra $L = \bigoplus_{n \in \mathbb{Z}} L_n$ induced by

defining, for all $v \in E^0$ and $e \in E^1$, $\deg(v) = 0$, $\deg(e) = 1$, $\deg(e^*) = -1$. Further, for each $n \in \mathbb{Z}$, the homogeneous component L_n is given by

$$L_n = \{ \sum k_i \alpha_i \beta_i^* \in L : |\alpha_i| - |\beta_i| = n \}.$$

If $p = e_1 \cdots e_n e_{n+1} \cdots$ is an infinite path in E , we follow Chen [14] to define, for each $n \geq 1$, $\tau^{\leq n}(p) = e_1 \cdots e_n$ and $\tau^{>n}(p) = e_{n+1} e_{n+2} \cdots$. Two infinite paths p, q are said to be *tail-equivalent* if there are positive integers m, n such that $\tau^{>m}(p) = \tau^{>n}(q)$. This defines an equivalence relation among the infinite paths in E and the equivalence class containing the path p is denoted by $[p]$. An infinite path p is said to be a *rational path* if it is tail equivalent to an infinite path $q = ccc \cdots$, where c is a closed path. An infinite path which is not rational is called an *irrational path*.

Convention: In what follows E denotes an arbitrary graph, K denotes a field and L denotes the Leavitt path algebra $L_K(E)$. Also, all the L -modules are assumed to be left L -modules.

2 Minimal Graded Left Ideals and the Graded Socle

The minimal left (right) ideals of Leavitt path algebras of arbitrary graphs were determined in [12]. It was proved that Lv is a minimal left ideal if and only if v is a line point. This can follow from a more general statement in the settings

of semiprime rings R . Namely, for an idempotent ϵ , $R\epsilon$ is a minimal left ideal if and only if $\epsilon R\epsilon$ is a division ring. In this section, we will use the graded version of the above statement to determine the graded minimal left (right) ideals. Specifically, we introduce the concept of a Laurent vertex in a graph and show that, in a Leavitt path algebra L , every minimal graded left ideal is isomorphic to $Lv(n)$ where v is either a Laurent vertex or a line point and $n \in \mathbb{Z}$. Using this, we obtain our structure theorem for the graded socle $\text{Soc}^{gr}(L)$ of a Leavitt path algebra L showing that $\text{Soc}^{gr}(L)$ is a ring direct sum of matrix rings of arbitrary size over K and/or over $K[x^t, x^{-t}]$ with appropriate grading, where $t \in \mathbb{N}$.

Recall that a graded L -module M is said to be graded-simple, if 0 and M are the only graded submodules of M . Observe that a graded-simple L -module need not be simple. For example, $M = K[x, x^{-1}]$ is graded-simple as a $K[x, x^{-1}]$ -module. But it is not simple, since there are infinitely many ideals of $K[x, x^{-1}]$ giving rise to proper submodules of M . On the other hand, a simple module need not be graded-simple. Indeed, if I is the ideal generated by any irreducible polynomial $p(x) \in K[x, x^{-1}]$, then $K[x, x^{-1}]/I$ is a simple module over $K[x, x^{-1}]$, but it is not graded-simple, as it is not even a graded module over $K[x, x^{-1}]$. We shall see in the sequel more occurrences of such examples.

We first start with a few preparatory lemmas.

Lemma 2.1 *Let I be a graded left ideal of a $\mathbb{Z}$ -graded ring R . Then I is a minimal graded left ideal if and only if $I(n)$ is a minimal graded left R -module for any $n \in \mathbb{Z}$.*

The proof is left to the reader.

Lemma 2.2 *If a vertex v is connected to a vertex u by a path p with no bifurcations, then $Lv(-|p|) \cong_{gr} Lu$ as graded left modules.*

Proof. Since $pp^* = v$ and $p^*p = u$, the map $av \mapsto ap$ is a graded isomorphism from $Lv(-|p|)$ to Lu . ■

The next Lemma was proved for non-graded simple left ideals in [12].

Lemma 2.3 *If, for a vertex v , Lv is a minimal graded left ideal of L , then $T_E(v)$ does not contain any bifurcating vertices.*

Proof. Indeed if $T_E(v)$ contains bifurcating vertices, choose a bifurcating vertex u such that there are no bifurcations in the path p connecting v to u . By Lemma 2.2, $Lu \cong Lv(-|p|)$ and by Lemma 2.1, Lu is a minimal graded left ideal. Let $e \neq f$ be two edges with $s(e) = u = s(f)$. Then Lee^* is a non-zero proper graded submodule of Lu contradicting the fact that Lu is a minimal graded left ideal. ■

Corollary 2.4 *If v is a vertex such that Lv is a minimal graded left ideal, then $|s^{-1}(v)| \leq 1$.*

The following Lemma will be used in the characterization of minimal graded left ideals of Leavitt path algebras. This is a graded version of a well-known result on semiprime rings. We will provide a simple proof for the graded version. Recall that a graded ring R is called graded semiprime if it has no non-zero nilpotent graded ideals.

Lemma 2.5 *Let Γ be a group, R a semiprime Γ -graded ring and ϵ be a homogeneous idempotent. Then $\epsilon R \epsilon$ is a graded division ring if and only if $R \epsilon$ (likewise, ϵR) is a minimal graded left (right) ideal. If $\epsilon R \epsilon$ is a field, then $R \epsilon$ (ϵR) is a minimal left (right) ideal.*

Proof. Suppose $\epsilon R \epsilon = \bigoplus_{\gamma \in \Gamma} \epsilon R_{\gamma} \epsilon$ is a graded division ring. Let $0 \neq a \in R \epsilon$ be a homogeneous element. It is enough if we show that $Ra = R \epsilon$. Now $aR \neq 0$, as $a = a\epsilon \in aR$. Since R is semiprime $(aR)^2 \neq 0$. Hence $aba \neq 0$ for some homogeneous element b . As $aba = a\epsilon b a \epsilon$, $\epsilon b a \epsilon \neq 0$ is a homogeneous element of the graded division ring $\epsilon R \epsilon$ with identity ϵ . Consequently, there is a homogeneous element y in $\epsilon R \epsilon$ such that $y\epsilon b a \epsilon = \epsilon$. Then for any $x \in R \epsilon$, we have $x = x\epsilon = xy\epsilon b a \epsilon \in Ra$. Hence $R \epsilon = Ra$, thus proving $R \epsilon$ is a minimal graded left ideal.

Conversely, suppose $R \epsilon$ is a minimal graded left ideal. Let $\epsilon x \epsilon$ be a homogeneous element of degree n in $\epsilon R \epsilon$. Clearly the right multiplication by $\epsilon x \epsilon$ is a non-zero graded morphism $\theta : R \epsilon(-n) \rightarrow R \epsilon$. Since $R \epsilon$ is a graded minimal left ideal, θ is actually a graded isomorphism. Let $\theta^{-1}(\epsilon) = \epsilon y \epsilon$. Then for any $a \in R \epsilon$ we have $\theta^{-1}(a) = \theta^{-1}(a\epsilon) = a\theta^{-1}(\epsilon) = a\epsilon y \epsilon$. Consequently, $\epsilon = \theta^{-1}\theta(\epsilon) = (\epsilon x \epsilon)(\epsilon y \epsilon)$. Similarly, $\epsilon = \theta\theta^{-1}(\epsilon) = (\epsilon y \epsilon)(\epsilon x \epsilon)$. Hence $\epsilon x \epsilon$ is invertible and we conclude that $\epsilon R \epsilon$ is a graded division ring. ■

Definition 2.6 *A vertex v is called a Laurent vertex if $T_E(v)$ consists of the set of all vertices on a single path $\gamma = \mu c$ where μ is a path without bifurcations starting at v and c is a cycle without exits based on a vertex $u = r(\mu)$.*

Lemma 2.7 *For any $v \in E^0$, the following hold:*

- (1) *If v is a line point, then $vLv \cong_{gr} K$.*
- (2) *If v is a Laurent vertex, then $vLv \cong_{gr} K[x^n, x^{-n}]$ for some integer $n \geq 1$.*

Proof. Now the statement (1) is well-known without the isomorphism being a graded isomorphism (see e.g. Proposition 2.7, [12]). Here is a short proof to get a graded version of this statement. We first show that if p and q are paths such that $v = s(p) = s(q)$, then $q = pt$ or $p = qs$, where s, t are paths in E . Suppose first the $|q| \geq |p|$. If $|p| = 0$, then $p = v$ and $q = vq$. We proceed by induction on $|p|$. Suppose the statement holds when $|p| = n \geq 0$. Let $|p| = n + 1$. Write $p = ep'$ and $q = fq'$ where $e, f \in E^1$. Since v is a line point, $e = f$ and $r(e)$ is also a line point. By induction, $q' = p't$. Then $q = fq' = ep't = pt$. The case when $|p| > |q|$ is similar.

Now consider the non-zero monomial vpq^*v . Note that $s(p) = s(q) = v$. If $|q| \geq |p|$, then $q = pt$ by the above argument. Thus $vpq^*v = vpt^*p^*v$. It follows

that $s(t) = r(t)$. Since v is a line point, t has to be a vertex, namely, $r(p)$. Then $vpq^*v = vpt^*p^*v = v$. The case when $|p| > |q|$ is similar. Thus $vLv \cong_{gr} K$ where K is a graded ring concentrated in degree 0.

(2) Since v is a Laurent vertex, $T_E(v)$ consists of a single path μc where μ is a path without bifurcations starting at v and ending at u and, c is a cycle of length n without exits based at u . By Lemma 2.2, $Lv(-n) \cong_{gr} Lu$. So

$$vLv \cong_{gr} \text{End}(Lv(-n)) \cong_{gr} \text{End}(Lu) \cong_{gr} uLu$$

. First observe that, since c is a cycle without exits, $cc^* = u = c^*c$ and any non-zero term upq^*u , where p and q are paths, simplifies to an integer power of c or c^* . Consequently, if $a = \sum_i uk_i p_i q_i^* u$ is a non-zero element of uLu , then a is a K -linear sum of powers of c and c^* . Also note that in the graded subring uLu , $\deg(c) = n$, the length of c and $\deg(c^*) = -n$. Then the map sending u to 1, c to x^n and c^* to x^{-n} defines a graded isomorphism of uLu to $K[x^n, x^{-n}]$ whose grading is induced by that of $K[x, x^{-1}]$. ■

Proposition 2.8 *Let $v \in E^0$. Then Lv is a minimal graded left ideal if and only if v is either a line point or a Laurent vertex.*

Proof. Suppose Lv is a minimal graded left ideal. Assume v is not a line point. Then $T_E(v)$ contains a vertex u which is either a bifurcating vertex or is the base of a cycle. By Lemma 2.3, $T_E(v)$ contains no bifurcations and so u is the base of a cycle c which obviously has no exits. In this case, $T_E(v)$ consists of a single path $\gamma = \mu c$ where μ is a path without bifurcations with $v = s(\mu)$ and $u = r(\mu) = s(c)$. Thus v is a Laurent vertex.

Conversely, suppose v is a line point. Then, by Lemma 2.7, $vLv \cong_{gr} K$ and so Lv is a minimal left ideal by Lemma 2.5. Lv is also a graded left ideal since v is a homogeneous idempotent of degree 0. Suppose, on the other hand, v is a Laurent vertex. By Lemma 2.7, vLv is graded isomorphic to the graded division ring $K[x^n, x^{-n}]$, where $n \geq 1$. By Lemma 2.5, Lv is a minimal graded left ideal. ■

Proposition 2.9 *A graded left ideal S of L is a minimal graded left ideal of L if and only if $S = La$ where a is a homogeneous element and $La \cong_{gr} Lu(n)$, where u is either a line point or a Laurent vertex and $n \in \mathbb{Z}$.*

Proof. Suppose $S = La$ is a minimal graded left ideal. We can assume that a is a homogeneous element. This is because, if $a = a_1 + \cdots + a_m$ where the a_i are homogeneous, then each $a_i \in S$ as S is graded. Moreover $La_i = S$ by graded-simplicity. So write $S = La$ for some homogeneous element a . By (Proposition 3.1, [12]), there exist paths μ, ν such that $\mu^* a \nu = kw$ or a polynomial $f(c, c^*)$, where w is a vertex, $k \in K$, $f(x, x^{-1}) \in K[x, x^{-1}]$ and c is a cycle without exits. Consider the case when $\mu^* a \nu = f(c, c^*)$. Since $\mu^* a \nu$ is homogeneous monomial, $\mu^* a \nu = k_i c^{n_i} (c^*)^{m_i}$ where $k_i \in K$ and m_i, n_i are integers. Then $\gamma^* a \delta = k' u$ where $\gamma = \mu c^{n_i}$, $\delta = \nu c^{m_i}$ and $k' \in K$. Thus, in either case, for suitable paths

α, β with β having no bifurcations, we get that $\alpha^*a\beta = ku$ where u is a vertex and $k \in K$. Now the map $\phi : La(-|\beta|) \rightarrow Lu$ given by $\phi(xa) = xa\beta$ is a non-zero graded morphism and is indeed a graded isomorphism by the simplicity of La . Thus $La(-|\beta|) \cong_{gr} Lu$. By Proposition 2.8, u is either a line point or a Laurent vertex. ■

From the preceding results, we obtain a description of the graded socle of a Leavitt path algebra L . Recall that for a ring R , $Soc^{gr}(R)$ denotes the sum of all minimal graded left ideals of R . Similar to the non-graded case, for semi-prime graded rings R (such as Leavitt path algebras L), the left and the right graded socles coincide. Even though $Soc(L)$ is a graded ideal, it need not be equal to $Soc^{gr}(L)$, as is clear from considering $L = K[x, x^{-1}]$. However, as noted in [24], $Soc(R) \subseteq Soc^{gr}(R)$, for any graded ring R . For a Leavitt path algebra L , one can also derive this from Proposition 2.8 and the fact that $Soc(L)$ is the sum of all the left ideals Lv where v is a line point [12].

Grading of a matrix ring over a $\mathbb{Z}$ -graded ring: In preparation of the structure theorem on graded socle of L and also for use in Section 6, we wish to recall the grading of matrices of finite order and then indicate how to extend this to the case of infinite matrices in which at most finitely many entries are non-zero (see [20] and [24]).

Let Γ be an additive abelian group, A be a Γ -graded ring and $(\delta_1, \dots, \delta_n)$ an n -tuple where $\delta_i \in \Gamma$. Then $M_n(A)$ is a Γ -graded ring with the λ -homogeneous components of $n \times n$ matrices

$$M_n(A)(\delta_1, \dots, \delta_n)_\lambda = \begin{pmatrix} A_{\lambda+\delta_1-\delta_1} & A_{\lambda+\delta_2-\delta_1} & \cdot & \cdot & \cdot & A_{\lambda+\delta_n-\delta_1} \\ A_{\lambda+\delta_1-\delta_2} & A_{\lambda+\delta_2-\delta_2} & & & & A_{\lambda+\delta_n-\delta_2} \\ & & & & & \\ & & & & & \\ A_{\lambda+\delta_1-\delta_n} & A_{\lambda+\delta_2-\delta_n} & & & & A_{\lambda+\delta_n-\delta_n} \end{pmatrix}. \quad (1)$$

This shows that for each homogeneous element $x \in A$,

$$\deg(e_{ij}(x)) = \deg(x) + \delta_i - \delta_j, \quad (2)$$

where $e_{ij}(x)$ is a matrix with x in the ij -position and with every other entry 0.

Now let A be a Γ -graded ring and let I be an arbitrary infinite index set. Denote by $M_I(A)$ the matrix with entries indexed by $I \times I$ having all except finitely many entries non-zero and for each $(i, j) \in I \times I$, the ij -position is denoted by $e_{ij}(a)$ where $a \in A$. Considering a "vector" $\bar{\delta} := (\delta_i)_{i \in I}$ where $\delta_i \in \Gamma$ and following the usual grading on the matrix ring (see (1),(2)), define, for each homogeneous element a ,

$$\deg(e_{ij}(a)) = \deg(a) + \delta_i - \delta_j. \quad (3)$$

This makes $M_I(A)$ a Γ -graded ring, which we denote by $M_I(A)(\bar{\delta})$. Clearly, if I is finite with $|I| = n$, then the graded ring coincides (after a suitable permutation) with $M_n(A)(\delta_1, \dots, \delta_n)$.

Suppose E is a finite acyclic graph consisting of exactly one sink v . Let $\{p_i : 1 \leq i \leq n\}$ be the set of all paths ending at v . Then it was shown in (Lemma 3.4, [3])

$$L_K(E) \cong M_n(K) \quad (4)$$

under the map $p_i p_j^* \mapsto e_{ij}$. Now taking into account the grading of $M_n(K)$, it was further shown in (Theorem 4.14, [20]) that the same map induces a graded isomorphism

$$L_K(E) \longrightarrow M_n(K)(|p_1|, \dots, |p_n|) \quad (5)$$

$$p_i p_j^* \mapsto e_{ij}.$$

In the case of a comet graph E (that is, a finite graph E in which every path eventually ends at a vertex on a cycle c without exits), it was shown in [4] that the map

$$L_K(E) \longrightarrow M_n(K[x, x^{-1}]) \quad (6)$$

$$p_i c^k p_j^* \mapsto e_{ij}(x^k)$$

induces an isomorphism. Again taking into account the grading, it was shown in (Theorem 4.20, [20]) that the map

$$L_K(E) \longrightarrow M_n(K[x^{|c|}, x^{-|c|}](|p_1|, \dots, |p_n|)) \quad (7)$$

$$p_i c^k p_j^* \mapsto e_{ij}(x^{k|c|}).$$

induces a graded isomorphism. Later in the paper [2], the isomorphisms (4) and (6) were extended to infinite acyclic and infinite comet graphs respectively (see Proposition 3.6 [2]). The same isomorphisms with the grading adjustments will induce graded isomorphisms for Leavitt path algebras of such graphs. We now describe this extension below.

Let E be a graph such that no cycle in E has an exit and such that every infinite path contains a line point or is tail-equivalent to a rational path $ccc \dots$ where c is a cycle (without exits). Define an equivalence relation in the set of all line points in E by setting $u \sim v$ if $T_E(u) \cap T_E(v) \neq \emptyset$. Let X be the set of representatives of distinct equivalence classes of line points in E , so that for any two line points $u, v \in X$ with $u \neq v$, $T_E(u) \cap T_E(v) = \emptyset$. For each vertex $v_i \in X$, let $\bar{p}^{v_i} := \{p_s^{v_i} : s \in \Lambda_i\}$ be the set of all paths that end at v_i , where Λ_i is an index set which could possibly be infinite. Denote by $|\bar{p}^{v_i}| = \{p_s^{v_i} : s \in \Lambda_i\}$.

Let Y be the set of all distinct cycles in E . As before, for each cycle $c_j \in Y$ based at a vertex w_j , let $\bar{q}^{w_j} := \{q_r^{w_j} : r \in \Upsilon_j\}$ be the set of all paths that end at w_j that do not include all the edges of c_j where Υ_j is an index set which could possibly be infinite. Let $|\bar{q}^{w_j}| := \{q_r^{w_j} : r \in \Upsilon_j\}$. Then the isomorphisms (5) and (7) extend to a $\mathbb{Z}$ -graded isomorphism

$$L_K(E) \cong_{gr} \bigoplus_{v_i \in X} M_{\Lambda_i}(K)(|\bar{p}^{v_i}|) \oplus \bigoplus_{w_j \in Y} M_{\Upsilon_j}(K[x^{|c_j|}, x^{-|c_j|}](|\bar{q}^{w_j}|)) \quad (8)$$

where the grading is as in (3).

We are now ready to prove the main theorem of this section.

Theorem 2.10 *Let E be an arbitrary graph and $L = L_K(E)$. Then the graded socle $Soc^{gr}(L)$ is the two-sided ideal generated by the set consisting of all the line points and all the Laurent vertices in E and*

$$Soc(L)^{gr} \cong_{gr} \bigoplus_{v_i \in X} M_{\Lambda_i}(K)(|p^{v_i}|) \oplus \bigoplus_{w_j \in Y} M_{\Upsilon_j}(K[x^{t_j}, x^{-t_j}])(|q^{w_j}|)$$

where Λ_i, Υ_j are suitable index sets, the t_j are positive integers, X is the set of representatives of distinct equivalence classes of line points in E and Y is the set of all distinct cycles (without exits) in E .

Proof. Let I be the two-sided ideal of L generated by all the line points and all the Laurent vertices in E . Now $I \subseteq Soc^{gr}(L)$ since, by Proposition 2.8, $Lu \subset Soc^{gr}(L)$ if u is a line point or a Laurent vertex. To prove the reverse inclusion, suppose $S = La$ is a minimal graded left ideal of L , where a is a homogeneous element. It is clear from Proposition 2.9 and its proof, that there is a vertex v which is either a line point or a Laurent vertex and a path β without exits such that $La(-|\beta|) = Lv\beta^*$. Consequently, $La(-|\beta|) \subset I$. Since I is a graded ideal, $S = La \subset I$. This proves that $I = Soc^{gr}(L)$. Let A be the two-sided ideal generated by the set H of all the line points in E and B be the two-sided ideal generated by the set H' of all the Laurent vertices in E . Clearly, $H \cap H' = \emptyset$. This implies that the intersection $\bar{H} \cap \bar{H}'$ of the saturated closures of H, H' is empty (See Lemma 1.4, [13]). Hence $A \cap B = 0$. So $Soc^{gr}(L) = A \oplus B$. By Theorem 3.7 and Propositions 3.5 and 3.6 of [2], there are algebraic isomorphisms $A \cong \bigoplus_{i \in I} M_{\Lambda_i}(K)$ and $B \cong \bigoplus_{j \in J} M_{\Upsilon_j}(K[x^{t_j}, x^{-t_j}])$

where Λ_i, Υ_j are arbitrary index sets and t_j are integers ≥ 1 . Now A and B are graded ideals of L and the indicated isomorphisms induce the graded isomorphism stated in (8) above, thus showing that

$$Soc(L)^{gr} \cong_{gr} \bigoplus_{v_i \in X} M_{\Lambda_i}(K)(|p^{v_i}|) \oplus \bigoplus_{w_j \in Y} M_{\Upsilon_j}(K[x^{t_j}, x^{-t_j}])(|q^{w_j}|).$$

■

Remark: In (Theorem 3.9, [2]), it is proved that a Leavitt path algebra L is categorically noetherian (that is, submodules of finitely generated left/right L -modules are finitely generated) exactly when $L \cong \bigoplus_{i \in I} M_{\Lambda_i}(K) \oplus \bigoplus_{j \in J} M_{\Lambda_j}(K[x, x])$

where Λ_i, Λ_j are suitable index sets and the t_j are positive integers. In view of Theorem 2.10, we conclude that a Leavitt path algebra L over an arbitrary graph is categorically noetherian if and only if L coincides with its graded socle.

3 Graded E-Algebraic Branching Systems and Graded Simple Modules

Given an arbitrary graph E , the concept of an E -algebraic branching system was studied by Gonçalves and Royer in [18] and was used to provide representations for a Leavitt path algebra L of the graph E . In this section, we define a grading on these algebraic branching systems and use the graded systems to construct various graded-simple modules over L . We also describe the corresponding annihilating ideals of these graded-simple L -modules. It may be of some interest to note that, unlike the ungraded case, the annihilator ideal of a graded-simple L -module need not be primitive, but it is always a graded prime ideal of L . Conversely, when E is row-finite or E^0 is countable, we show that every graded prime ideal of L which is not primitive occurs as the annihilator of some graded-simple L -module which is not simple. Using infinite paths and sinks in E , Chen [14] constructed simple L -modules and showed that these simple modules can also be constructed by using algebraic branching systems. This approach was further expanded in [11] and [26] to introduce new types of simple L -modules induced by vertices which are infinite emitters in E . All these new modules are now called Chen simple L -modules. Examples show that not all graded-simple L -modules are Chen simple modules and likewise, not all Chen simple L -modules are graded-simple. We show that the Chen simple module $V_{[p]}$ corresponding to an infinite path p is graded-simple if and only if p is an infinite irrational path in E .

Definition 3.1 *Let E be an arbitrary graph. An E -algebraic branching system consists of a set X and a family of its subsets $\{X_v, X_e : v \in E^0, e \in E^1\}$ such that*

- (1) $X_v \cap X_w = \emptyset = X_e \cap X_f$ for $v, w \in E^0$ with $v \neq w$ and $e, f \in E^1$ with $e \neq f$;
- (2) $X_e \subseteq X_{s(e)}$ for $e \in E^1$;
- (3) For all $v \in E^0$, $X_v = \bigcup_{e \in s^{-1}(v)} X_e$;
- (4) For each $e \in E^1$, there exists a bijection $\sigma_e : X_{r(e)} \rightarrow X_e$.

For our purposes, we also assume that X is saturated, that is, $X = \bigcup_{v \in E^0} X_v$.

Definition 3.2 *The E -branching system is called graded if there is a map $\deg : X \rightarrow \mathbb{Z}$ such that*

- (5) $\deg(\sigma_e(x)) = \deg(x) + 1$.

Given an E -algebraic branching system X , we define a left module over $L := L_K(E)$ as follows: Consider the K -vector space $M(X)$ having X as a basis. Following [18], we define, for each vertex v and each edge e in E , linear transformations P_v, S_e and S_{e^*} on $M(X)$ as follows:

Definition 3.3 For all paths $x \in X$,

$$\begin{aligned} (I) \quad P_v(x) &= \begin{cases} x, & \text{if } x \in X_v \\ 0, & \text{otherwise} \end{cases} \\ (II) \quad S_e(x) &= \begin{cases} \sigma_e(x), & \text{if } x \in X_{r(e)} \\ 0, & \text{otherwise} \end{cases} \\ (III) \quad S_{e^*}(x) &= \begin{cases} \sigma_e^{-1}(x), & \text{if } x \in X_e \\ 0, & \text{otherwise} \end{cases} \end{aligned}$$

Then it is straightforward to check that the endomorphisms $\{P_u, S_e, S_{e^*} : u \in E^0, e \in E^1\}$ satisfy the defining relations (1) - (4) of the Leavitt path algebra L (see [18]). This induces an algebra homomorphism ϕ from L to $\text{End}_K(M(X))$ mapping u to P_u , e to S_e and e^* to S_{e^*} ([18], Theorem 2.2). Then $M(X)$ can be made a left module over L via the homomorphism ϕ . We denote this L -module operation on $M(X)$ by $\cdot$.

If X is a graded branching system, then define, for each $i \in \mathbb{Z}$, the homogeneous component

$$M(X)_i = \left\{ \sum_{x \in X} k_x x \in M(X) : \deg(x) = i \right\}.$$

It is easy to see that

$$M(X) = \bigoplus_{i \in \mathbb{Z}} M(X)_i$$

and that $M(X)$ is a $\mathbb{Z}$ -graded left L -module.

We shall now provide three illustrations of constructing graded-simple L -modules using appropriate E -algebraic branching systems.

A) A graded-simple module which is not simple: Let E be a graph. Let $u \in E^0$ be a Laurent vertex so that $T_E(u)$ consists of a single path $\gamma = \mu c$ where γ is a path without bifurcations and c is a cycle without exits based at a vertex v . Let $X = \{pq^* : p, q \text{ paths with } r(q^*) = v\}$. We make X a graded branching system as follows:

$$\begin{aligned} X_w &= \{pq^* \in X : s(p) = w\}, \text{ where } w \in E^0; \\ X_e &= \{pq^* \in X : e \text{ initial edge of } p\}, \text{ where } e \in E^1; \\ \sigma_e : X_{r(e)} &\rightarrow X_e \text{ given by } pq^* \mapsto epq^*, \text{ a bijection;} \\ \deg : X &\rightarrow \mathbb{Z}, \text{ given by } pq^* \mapsto |pq^*| = |p| - |q|. \end{aligned}$$

The only non-trivial condition of the algebraic branching that needs to be checked here is $X_w = \bigcup_{e \in s^{-1}(w)} X_e$. Let $pq^* \in X_w$. If $|p| \geq 1$ then clearly $p = et$,

where $s(e) = w$. Thus $pq^* = etq^* \in X_e$. If $|p| = 0$, then $pq^* = q^*$ such that $s(q^*) = r(q) = w$ and, by hypothesis on X , $r(q^*) = s(q) = v$. Since $\gamma = \mu c$ has no exits, the path q is part of γ and w will emit only one edge e . Then $q^* = ee^*q^* \in X_e$ thus showing $X_w \subseteq \bigcup_{e \in s^{-1}(w)} X_e$. The converse inclusion is

obvious.

Then conditions (I),(II),(III) make $M(X)$ a graded L -module which we denote by N_{vc} .

Next we wish to list some of the properties of module N_{vc} in Theorem 3.5. Its proof needs the following descriptions of the graded prime and graded primitive ideals of a Leavitt path algebra L of an arbitrary graph E stated in Theorems 3.12 and 4.3 of [25]. Recall ([5]), subset X of E^0 is said satisfy the *countable separation property* if there is a countable subset Y of E^0 with the property that to each $u \in X$, there is a $v \in Y$ such that $u \geq v$.

Theorem 3.4 ([25]): *Let P be an ideal of L with $P \cap E^0 = H$. Then*

(a) *P is a graded prime ideal if and only if either (i) $P = I(H, B_H)$ (where B_H may be empty) with $E^0 \setminus H$ downward directed or (ii) $P = I(H, B_H \setminus \{u\})$ for some $u \in B_H$ and $w \geq u$ for every $w \in E^0 \setminus H$.*

(b) *P is a graded primitive ideal if and only if either (i) P is a graded prime ideal of the form $I(H, B_H)$ such that $E \setminus H$ satisfies Condition (L) and the countable separation property or (ii) P is the prime ideal of type (ii) described above.*

The next theorem lists some properties of the module N_{vc} .

Theorem 3.5 *Let v be a Laurent vertex based at a cycle c . Then we have the following:*

(1) *The module N_{vc} is a graded-simple L -module but is not a simple L -module;*

(2) *The annihilator ideal of N_{vc} is the ideal $I(H(v), \emptyset)$ where $H(v) = \{u \in E^0 : u \not\geq v\}$. It is a graded prime ideal of L which is not primitive;*

(3) *Conversely, if E is row-finite or if E^0 is countable, then every graded prime ideal P of L which is not primitive is the annihilator of a graded-simple module which is not simple.*

Proof. (1) Suppose U is a non-zero graded submodule of N_{vc} . Let a be a

non-zero homogeneous element in U , say, $a = \sum_{i=1}^n k_i p_i q_i^*$, where, by definition,

$r(q_i^*) = v$. Multiplying by p_1^* on the left, and replacing a by $p_1^* a$, we may assume that the first term of a is $k_1 q_1^*$. Again multiplying on the left by $s(q_1^*)$ and gathering all the non-zero terms, we may assume without loss of generality that

$$a = k_1 q_1^* + k_2 p_2 q_2^* + \cdots + k_r p_r q_r^*$$

where $|q_1| > 0$, $s(p_i) = s(q_1^*) = r(q_1)$ and $s(q_1) = r(q_1^*) = r(q_i^*) = s(q_i) = v$ for all i . Since the path c has no exits, q_1 lies entirely on c and this implies that the p_i lie entirely on c . Further the q_i also lie on c with $s(q_i) = v$ and $r(q_i) = r(p_i)$. Since $\deg(p_i q_i^*) = \deg(q_1^*)$ for all i , we conclude that each $p_i q_i^* = q_1^*$. So $a = k q_1^*$ for some $k \in K$. Then $k^{-1} q_1 a = v \in U$ and so $U = N_{v\infty}$. Thus $N_{v\infty}$ is a graded-simple L -module. But N_{vc} is not a simple L -module. Because, if f is an edge with $r(f) = v$, then it can be readily seen that $v \notin L(v + f)$, thus showing that $(v + f)$ generates a non-zero proper submodule of N_{vc} .

(2) First observe that the annihilator Q of a graded simple module S is always a graded prime ideal: Q is graded, because if $a \in Q$ with $a = a_1 + \cdots + a_n$ a graded sum of homogeneous elements a_i , then for any homogeneous element $x \in S$, $0 = ax = a_1x + \cdots + a_nx$ implies $a_ix = 0$ for all i . Hence each $a_i \in Q$. To see that Q is prime, suppose A, B are graded ideals such that $A \not\subseteq Q$ and $B \not\subseteq Q$. Then AS is a non-zero graded submodule of S and so $AS = S$, by simplicity. Similarly $BS = S$. Then $ABS = AS = S$. Hence $AB \not\subseteq Q$, thus proving that Q is graded prime. Thus the annihilator P of N_{vc} must be a graded prime ideal of L . From the description of N_{vc} it is clear that $H(v) = \{u \in E^0 : u \not\geq v\} \subset P$. Actually, $P \cap E^0 = H(v)$, since $w \geq v$ for all $w \in E^0 \setminus H(v)$. Also, since $B_H \subseteq E^0 \setminus H$, we have for each $u \in B_H$ a path p connecting u to v . Clearly then

$$u^{H(v)}p = (u - \sum_{e \in s^{-1}(u), r(e) \in H(v)} ee^*)p = p \neq 0.$$

So $u^{H(v)} \notin P$. It then follows from Theorem 3.4(a) that $P = I(H(v), \emptyset)$. As $E \setminus H(v)$ contains the cycle c without exits, it does not satisfy Condition (L) and hence by Theorem 3.4(b) that P is not a primitive ideal of L .

(3) Suppose P is a graded prime ideal of L which is not primitive. Let $H = P \cap E^0$. By Theorem 3.4, $E^0 \setminus H$ is downward directed and $P = I(H, B_H)$. Also, $E \setminus H$ satisfies the countable separation property whenever $E \setminus H$ is row-finite or $E^0 \setminus H$ is countably infinite [5]. Then the hypothesis that P is not primitive implies, by Theorem 3.4, that $E \setminus H$ does not satisfy Condition (L). Hence there is a cycle c based at a vertex v and without exits in $E \setminus (H, B_H)$. Thus v is a Laurent vertex in $E \setminus H$. The denoting $\bar{v} = v + P$, we appeal to Proposition 2.8 to conclude that $S = (L/P)\bar{v} \cong L_K(E \setminus (H, B_H))v$ is a graded-simple L/P -module which obviously not a simple L/P -module (as v is not a line point in $E \setminus (H, B_H)$). Since $w \geq v$ for every vertex $w \in E \setminus (H, B_H)$, it is then clear that the annihilator of S as an L/P -module is $\{0\}$. Consequently, as an L -module S is a graded-simple module which is not simple and its annihilator ideal is P . ■

B) A graded-simple module which is also simple: Let E be any graph and let $v \in E^0$ be an infinite emitter. Let X be the set of all the finite paths ending at v . We make X a graded branching system as follows:

$$X_w = \{q \in X : s(q) = w\}, \text{ where } w \in E^0;$$

$$X_e = \{q \in X : e \text{ is the initial edge of } q\}, \text{ where } e \in E^1;$$

$$\text{A bijection } \sigma_e : X_{r(e)} \rightarrow X_e \text{ given by } \sigma_e(q) = eq;$$

$$\deg : X \rightarrow \mathbb{Z} \text{ is given by } \deg(p) = |p|.$$

Then the K -vector space $M(X)$ having X as a basis becomes a left L -module by using the condition (I),(II),(III). It becomes a graded module by defining, for each $i \in \mathbb{Z}$,

$$M(X)_i = \left\{ \sum_{p_i \in X} k_i p_i \in M(X) : |p_i| = i \right\}$$

and $M(X) = \bigoplus_{i \in \mathbb{Z}} M(X)_i$. Now the module $M(X)$ is the same thing as the module $S_{v\infty}$ introduced in [26], since the construction $M(X)$ is essentially a reformulation, using the E-algebraic branching systems, of the construction of the module $S_{v\infty}$. In (Proposition 2.3, [26]), it was shown that $S_{v\infty}$ becomes a simple L -module. Since $S_{v\infty} = M(X)$ is also graded, $S_{v\infty}$ is a graded-simple left L -module. As noted in [26], $S_{v\infty}$ gives rise to three distinct classes of non-isomorphic simple and hence graded-simple L -modules depending on the vertex v satisfying one of the following three properties, where $N(v) = \{u \in E : u \geq v\}$:

- (i) $|s^{-1}(v) \cap r^{-1}(N(v))| = 0$;
- (ii) $0 < |s^{-1}(v) \cap r^{-1}(N(v))| < \infty$;
- (iii) $|s^{-1}(v) \cap r^{-1}(N(v))|$ is infinite.

Remark: (i) Suppose w is a sink in E . Define X to be the set of all finite paths ending at w . Proceed as in paragraph B) above to make X a graded algebraic branching system and construct the graded L -module $M(X)$. This module was denoted by N_w in ([14]) where it was shown to be simple. It is thus a graded-simple L -module which is also simple.

(ii) The above approach to defining a graded simple module does not work if v is a regular vertex. This is because, in this case, the linear transformations P_v, S_e, S_{e^*} do not satisfy the CK-2 relation $\sum_{e \in s^{-1}(v)} S_e S_{e^*} = P_v$. Because on

the one hand $(\sum_{e \in s^{-1}(v)} S_e S_{e^*})(v) = 0$ but on the other hand $P_v(v) = v \neq 0$.

C) A simple module which is not graded-simple: Let E be any graph and let p be an infinite path in E . Define an E -algebraic branching system with $X = [p]$, the set of all infinite paths tail-equivalent to p .

For each $v \in E^0$, let $X_v = \{q \in [p] : s(q) = v\}$

For each $e \in E^1$, let $X_e = \{q \in [p] : e \text{ is the initial edge of } q\}$.

Define $\sigma_e : X_{r(e)} \rightarrow X_e$ by $q \mapsto eq$.

It was shown by Chen in ([14]) that $M(X)$ becomes a simple left L -module and denoted it by $V_{[p]}$.

Next we show that infinite rational paths give rise to non-gradable simple L -modules. We begin with a simple lemma.

Proposition 3.6 *Let E be an arbitrary graph and let p be an infinite path in E . Then the corresponding Chen simple left L -module $V_{[p]}$ is a graded-simple L -module if and only if p is an irrational infinite path in which case p is a homogeneous element of $V_{[p]}$.*

Proof. Assume p is a rational path, say $p = ccc \cdots$ where c is a cycle. Suppose, by way of contradiction, that $V_{[p]}$ becomes a graded module under the defined $L_K(E)$ -module operation. First we claim that p must be a homogeneous ele-

ment. Suppose, on the contrary,

$$p = \sum_{i=1}^n h_i \quad (++)$$

where the h_i are non-zero homogeneous elements in $V_{[p]}$ and $h_i \neq p$ for all i .

Since $h_i \in V_{[p]}$, write $h_i = \sum_{j=1}^{t_i} k_{ij} q_{ij}$ where $k_{ij} \in K$, $q_{ij} \in [p]$ and the paths q_{ij}

are distinct for all i, j . Choose an integer n such that the paths $p^{\leq n}$ and the $q_{ij}^{\leq n}$ are distinct for all i, j . Multiplying the equation $(++)$ on the left with $(c^*)^{-n}$, we then get $p = 0$, a contradiction. Thus p is homogeneous. Since $p = cp$, we then get

$$\deg(p) = \deg(cp) = \deg(p) + |c|,$$

which is a contradiction. Hence $V_{[p]}$ cannot be graded as an L -module.

Conversely, suppose p is an infinite irrational path in E . For each $q \in [p]$, define $\deg(q) = m - n$, if m is the smallest positive integer such that $\tau^{>m}(q) = \tau^{>n}(p)$ for some n . It is then easy to check that the simple left L -module $V_{[p]}$ becomes a graded L -module. The proof that the irrational path p is homogeneous in $V_{[p]}$ is the same as was given in the preceding paragraph where, instead of multiplying the equation $(\#)$ on the left by $(c^*)^{-n}$, multiply by $(p^{\leq n})^*$ to get a contradiction. ■

4 Leavitt Path Algebras over which every graded-simple module is finitely presented

Leavitt path algebras $L_K(E)$ over which every non-graded simple modules are finitely presented seem to have interesting properties. These were completely described in [11] when E is a finite graph and in [27] when E is an arbitrary graph. In this section, we provide a complete description of the Leavitt path algebra L of an arbitrary graph E over which every graded-simple is finitely presented.

First observe that if every simple L -module is finitely presented, then every graded-simple module need not be finitely presented. For example, let L be the algebraic Toeplitz algebra so that $L = L_K(E)$, with E consisting of two vertices v, w and two edges e, f such that $v = s(e) = r(e) = s(f)$ and $w = r(f)$. It was shown in (Example 4.5, [26]) that every simple L -module is finitely presented. This also follows from Theorem 1.1 of [11]. On the other hand, if I denotes the ideal generated by w , then I is a graded ideal and $L/I \cong K[x, x^{-1}]$ is a graded-

simple L -module. But L/I is not finitely presented since $I = Lw \oplus \bigoplus_{n=0}^{\infty} Lf^*(e^*)^n$ is infinitely generated. As we shall see later, the condition that all the graded-simple modules are finitely presented places considerable restriction on the nature of the Leavitt path algebra L .

In [6], the authors investigate when the Chen simple module $V_{[p]}$ corresponding to an infinite path p is finitely presented. One of their key results (see Theorem 2.20, [6]) states, in essence, that if p is an infinite irrational path no vertex on which is either an infinite emitter or a line point, then the corresponding Chen simple L -module $V_{[p]}$ is not finitely presented. But the proof of this important result involves lengthy computations and it depends on the additional requirement that the vertices on this path p must be finite emitters. Using Theorem 4.2 below, we shall show how to get rid of this additional requirement for the vertices on the path p and to provide a short proof of a more general result (Theorem 4.6).

We first consider a graded reformulation of Proposition 2 in [7] which was proved without any reference to the grading of the Leavitt path algebra and which is very useful in this and the next section. A careful inspection of the construction of the morphism in their proof shows that the stated homomorphism θ , mentioned in Theorem 4.1 below, is indeed a graded morphism whose image is a graded submodule of L . We reformulate this theorem reflecting the grading of L .

Theorem 4.1 ([7]) *Let E be an arbitrary graph. Then the Leavitt path algebra $L := L_K(E)$ is a directed union of graded subalgebras $B = A \oplus K\epsilon_1 \oplus \cdots \oplus K\epsilon_n$ where A is the image of a graded homomorphism θ from a Leavitt path algebra $L_K(F_B)$ with F_B a finite graph (depending on B), the ϵ_i are homogeneous orthogonal idempotents and $\oplus$ is a ring direct sum. Moreover, if E is acyclic, so is each graph F_B and in this case θ is a graded monomorphism.*

Proof. For the proof of the first statement, see [7]. The second statement that θ is a monomorphism follows from the fact that, when E is acyclic, the mentioned finite graph F_B (being acyclic) satisfies Condition (L) and since θ does not vanish on the vertices of F_B , θ is a monomorphism by the graded uniqueness theorem. ■

Another useful result is the following theorem proved in [21]. This theorem was first proved for finite graphs E . Then it was extended to row-finite graphs and then to arbitrary graphs by the so-called desingularization process. However the desingularization of a graph does not always preserve the grading of the algebra. We show below how to fix this problem by using Theorem 4.1 and deriving a shorter proof of the general case when E is an arbitrary graph. Recall that a graded ring R is called *graded von Neumann regular* if for each homogeneous element x there is a homogeneous element y such that $xyx = x$.

Theorem 4.2 ([21]) *Every Leavitt path algebra L of an arbitrary graph E is a graded von Neumann regular ring.*

Proof. It was shown in (Theorem 10 (I), [21]) that if E is any finite graph, then L is graded von Neumann regular. Suppose now E is an arbitrary graph. By Theorem 4.1, L is a directed union of graded subalgebras $B = A \oplus K\epsilon_1 \oplus \cdots \oplus K\epsilon_n$ where A is the image of a graded homomorphism θ from a Leavitt path algebra $L_K(F_B)$ with F_B a finite graph (depending on B), the ϵ_i are orthogonal

idempotents and $\oplus$ is a ring direct sum. Since F_B is a finite graph, $L_K(F_B)$ and hence B is graded von Neumann regular by [21]. It is then clear from the definition that the directed union L is also graded von Neumann regular. ■

We summarize some of the properties of graded modules over graded rings without identity (such as the Leavitt path algebras) that we will be using. Even though a ring R without identity may not be projective as an R -module, it was shown in [13] that a Leavitt path algebra L is always projective as a left (right) L -module. Since L is also graded, it becomes a graded-projective L -module. Now L is graded von Neumann regular by Theorem 4.2. We shall be using the following graded version of a known result which can be obtained by a simple induction on the number of generators:

A finitely generated graded left ideal of L is a direct summand of L (**)

We also wish to note that, in spite of the fact that L may not have a multiplicative identity, maximal graded left ideals always exist in L . Because, if $v \in E^0$, then by Zorn's Lemma, Lv contains a maximal graded L -submodule N . Then $N \oplus \bigoplus_{u \in E^0, u \neq v} Lu$ is a maximal graded left ideal of L . Indeed, every

maximal graded left ideal M of L is of this form: Given M there is a unique vertex v such that $M = N \oplus \bigoplus_{u \in E^0, u \neq v} Lu$ where $N = Mv = M \cap Lv$ is a maximal graded L -submodule of Lv (see Lemma 3.1, Remark 3.2, [26]). Consequently every graded-simple L -module S of L is graded isomorphic to Lv/N for some vertex v and some maximal graded submodule N of Lv .

The following Lemma is well-known for non-graded modules over rings with identity. We will modify the proof since L need not have an identity and since we are dealing with graded modules.

Lemma 4.3 *If every graded-simple left L -module is projective, then L is graded semi-simple, that is, L is a direct sum of minimal graded left ideals.*

Proof. Our hypothesis implies that every maximal graded left ideal of L is a direct summand and so L contains minimal graded left ideals. By Zorn's Lemma choose a maximal family $\mathcal{F}$ of independent minimal graded left ideals of L and let $M = \bigoplus_{X \in \mathcal{F}} X$. We claim that $M = L$. Because, otherwise, there is a vertex $v \notin M$. In this case, we can embed (by Zorn's Lemma) $M \cap Lv$ inside a maximal graded L -submodule N of Lv which is easily seen as a direct summand of Lv , say $Lv = N \oplus X'$. Then $\mathcal{F} \cup \{X'\}$ contradicts the maximality of $\mathcal{F}$. Thus $L = \bigoplus_{X \in \mathcal{F}} X$ is a direct sum of minimal graded left ideals. ■

Before we state our next theorem, we consider the following preparatory lemma.

Lemma 4.4 *Let $v \in E^0$ be an infinite emitter. Then the corresponding graded-simple module $S_{v\infty}$ (considered in section 3) is not finitely presented.*

Proof. Suppose, on the contrary, $S_{v\infty}$ is finitely presented. Note that $S_{v\infty} \cong Lv/N$ for some N which will be finitely generated by supposition. Then, by statement (**) above, $S_{v\infty}$ will be a graded-projective L -module. This means that $S_{v\infty}$ is isomorphic to a minimal graded left ideal of L and hence, by Proposition 2.9, $S_{v\infty} \cong Lu(n)$ where u is either a line point or a Laurent vertex. Both cases lead to a contradiction. Because if u is a line point, then by Chen [14], $Lu \cong N_w$ or $V_{[p]}$ according as $T_E(u)$ is a finite path ending at a sink w or an infinite path p and this is in contradiction to Theorem 2.9 of [26] whose proof shows that $S_{v\infty}$ is not isomorphic to either N_w or to $V_{[p]}$. Likewise, if u is a Laurent vertex, then Lu , by Theorem 3.5 (1), cannot be a simple left ideal, but it was shown in [26] that $S_{v\infty}$ is indeed a simple left L -module. These contradictions show that $S_{v\infty}$ cannot be finitely presented. ■

Theorem 4.5 *Let E be an arbitrary graph. Then the following are equivalent for $L := L_K(E)$.*

- (1) *Every graded-simple left L -module is finitely presented;*
- (2) *L is graded semi-simple, that is, L is a direct sum of minimal graded left ideals;*
- (3) *The graph E is row-finite and for each vertex u there is an integer $n \geq 0$ such that every path beginning with u and having length $\geq n$ ends at a line point or a Laurent vertex.*

Proof. Assume (1). As we noted earlier, every graded-simple L -module S is graded isomorphic to Lv/N where v is a vertex and N is a graded submodule of Lv . Now L and hence Lv is a graded-projective L -module. So if S is finitely presented, then by Schanuel's Lemma, N becomes a finitely generated graded left ideal of L . By Theorem 4.2, L is graded von Neumann regular and so, by (**), N is a direct summand of L and that of Lv . This implies that S is graded-projective. Since every graded-simple L -module is graded-projective, we appeal to Lemma 4.3 to conclude that L is a graded semi-simple algebra. This proves (2).

Assume (2). Since every graded submodule of a graded semi-simple module is a direct summand and since L is graded projective, every graded homomorphic image of L and, in particular, every graded-simple L -module S is projective. Clearly S is finitely presented. This proves (1).

Assume (1) and (2). By Lemma 4.4, E must be row-finite. Also (2) implies $L = Soc^{gr}(L)$. From Theorem 2.10, we conclude that E^0 is the saturated closure of the set of all line points and Laurent vertices in E . This means (see e.g. Lemma 1.4, [8]) that for each vertex u there is an integer $n \geq 0$ such that every path beginning with u and having length $\geq n$ ends at a line point or a Laurent vertex, thus proving (3).

Assume (3). Since E is row-finite, the stated condition implies that E^0 is the saturated closure of the set of all line points and Laurent vertices in E . By Theorem 2.10, $L = Soc^{gr}(L)$ is a direct sum of minimal graded left ideals. This proves (2). ■

We next consider when the Chen simple module $V_{[p]}$ corresponding to an infinite path p is finitely presented.

As mentioned in the introductory paragraphs of this section, we now use Proposition 3.6, to present a short and simpler proof of a more general version of Theorem 2.20 of [6] without requiring the additional condition that no vertex on p is an infinite emitter.

Theorem 4.6 *Let E be an arbitrary graph, $L = L_K(E)$ and let p be an infinite irrational path in E . Then the following properties are equivalent:*

- (i) *The Chen simple left L -module $V_{[p]}$ is finitely presented;*
- (ii) *$V_{[p]}$ is a projective left L -module;*
- (iii) *The path p contains a vertex which is a line point.*

Proof. (i) $\Rightarrow$ (ii). As noted in Proposition 3.6, $V_{[p]}$ is a graded-simple L -module in which p is a homogeneous element. So if $v = s(p)$, we then get a graded exact sequence

$$0 \rightarrow N \rightarrow Lv \xrightarrow{\theta} V_{[p]} \rightarrow 0$$

where $\theta(av) = av \cdot p$. If $V_{[p]}$ is finitely presented, then N will be finitely generated. Since, by Theorem 4.2, L is graded von Neumann regular, statement (**) above implies that N will then be a direct summand of L . Thus the above graded exact sequence splits and we conclude that $V_{[p]}$ is graded-projective.

(ii) $\Rightarrow$ (iii). If $V_{[p]}$ is a projective left L -module, then, being simple, $V_{[p]}$ is isomorphic to a minimal left ideal Lu where u is a line point. By Proposition 4.3 of [14], p contains a vertex which is a line point, thus proving (iii).

(iii) $\Rightarrow$ (i). Again if p contains a vertex u which is a line point, then $V_{[p]} \cong Lu$, by ([14], Proposition 4.3). Since Lu is projective, $V_{[p]}$ is then clearly finitely presented. ■

5 Leavitt Path Algebras over which every simple module is graded

In this short section, we characterize the Leavitt path algebras of arbitrary graphs over which every simple left module is graded. A well-known theorem (see [29]) states that every two-sided ideal of a Leavitt path algebra $L_K(E)$ is graded if and only if the graph E satisfies Condition (K). A natural question is to investigate when every one-sided ideal of L is graded. We show that this happens exactly when the graph E contains no cycles. A useful role is played by Theorems 4.1 and 4.6.

Theorem 5.1 *Let E be an arbitrary graph. Then for the Leavitt path algebra $L := L_K(E)$ the following are equivalent:*

- (1) *Every left/right ideal of L is graded;*
- (2) *Every simple left/right L -module is graded;*
- (3) *The graph E is acyclic.*

Proof. (1) $\Rightarrow$ (2) Suppose S is a simple L -module. Since L is a ring with local units, there is a maximal left ideal M such that $S \cong L/M$. Now, by assumption,

M is a graded left ideal. Consequently, L/M and hence S can be made a graded L -module.

(2) $\Rightarrow$ (3). Now E must be acyclic, because otherwise E will contain a cycle c and corresponding to the infinite rational path $p = ccc \cdots$, the Chen simple module $V_{[p]}$ is, by Proposition 3.6, not a graded L -submodule, a contradiction. Thus E is acyclic.

(3) $\Rightarrow$ (1). Suppose E is acyclic. Now, by Theorem 4.1, L is a directed union of graded subalgebras B_λ where $\lambda \in I$, an index set and where each B_λ is a finite direct sum of copies of K and a graded homomorphic image of a Leavitt path algebra of a finite acyclic graph. By (Theorem 4.14, [20]), Leavitt path algebras of finite acyclic graphs are semi-simple algebras which have elementary gradings, that is, all the matrix units are homogeneous. Consequently, every left ideal of each B_λ is graded. Since the B_λ are graded subalgebras, each $B_\lambda = \bigoplus_{n \in \mathbb{Z}} (B_\lambda \cap L_n)$ where $L = \bigoplus_{n \in \mathbb{Z}} L_n$ is the $\mathbb{Z}$ -graded decomposition of L . Let M be a left ideal of L . To show that M is graded, we need only to show that $M = \bigoplus_{n \in \mathbb{Z}} (M \cap L_n)$. Let $a \in M$. Then, for some λ , $a \in M \cap B_\lambda$. Note that $M \cap B_\lambda = B_\lambda$ or a left ideal of B_λ . Since every left ideal of B_λ and in particular $M \cap B_\lambda$ is graded, we can write $a = a_{n_1} + \cdots + a_{n_k}$ where

$$a_{n_i} \in (M \cap B_\lambda) \cap (B_\lambda \cap L_{n_i}) \subseteq M \cap L_{n_i}$$

for $i = 1, \dots, k$. This shows that $M = \bigoplus_{n \in \mathbb{Z}} (M \cap L_n)$ and hence M is a graded left ideal of L ■

6 Graded self-injective Leavitt path algebras

It was shown in Theorem 4.2 that every Leavitt path algebra L is always graded von Neumann regular. On the other hand, a self-injective Leavitt path algebra is always von Neumann regular. So we wish to describe the subclass of graded self-injective Leavitt path algebras. The (non-graded) self-injective Leavitt path algebras were described in [13] as those associated to acyclic graphs. However, since the forgetful functor $U : GrA \rightarrow \text{mod } A$ is, in general, only a left-adjoint, this implies that the graded projective modules are projective, but the graded injective modules are not necessarily injective. In Theorem 6.7, we determine the class of graded self-injective Leavitt path algebras. These algebras are shown to coincide with their graded socle and thus are direct sums of matrix rings of arbitrary size over the field K and/or the ring $K[x^{t_i}, x^{-t_i}]$ endowed with a natural graded structure, where the t_i are integers.

We begin with the following Lemma which is folklore. This will be used in Lemma 6.2. For the sake of completeness, we outline the proof.

Lemma 6.1 *Let τ be an infinite cardinal and let K be any field. Then the K -dimension of the direct product $P = \prod_{\tau} K$ of τ copies of K is $\geq 2^\tau$.*

Proof. Clearly $|P| = \dim_K(P) \cdot |K|$. So if $|K| \leq \tau$, then $\dim_K(P) = |P| = 2^\tau$. Suppose $|K| > \tau$. Let Q be the prime subfield of K , so $|Q|$ is finite or countably infinite. If $P' = \prod_{\tau} Q$, then, as noted earlier, $\dim_Q(P') = 2^\tau$. Let $\bar{P} = K \otimes_Q P'$.

Then $\dim_K(\bar{P}) = 2^\tau$. Since $\bar{P} \subseteq P$, $\dim_K(P) \geq 2^\tau$. ■

A useful tool in describing the graded self-injective Leavitt path algebras is the following.

Lemma 6.2 *Suppose $v \in E^0$ such that vLv has K -dimension σ . If $\{Lx_i : i \in I\}$ is an independent family of graded submodules of Lv and $|I| = \tau \geq \max\{\sigma, \aleph_0\}$, then Lv is not graded injective as a left L -module.*

Proof. Suppose, by way of contradiction, Lv is a graded injective L -module. Let $S = \bigoplus_{i \in I} Lx_i$. Then from the graded exact sequence $0 \rightarrow S \rightarrow Lv$ we get the exact sequence

$$\text{Hom}_{gr}(Lv, Lv) \rightarrow \text{Hom}_{gr}(S, Lv) \rightarrow 0.$$

Now $\text{Hom}_{gr}(Lv, Lv) \cong vL_0v$ and

$$\text{Hom}_{gr}(S, Lv) \supseteq \text{Hom}_{gr}(S, S) \supseteq \prod_{i \in I} \text{Hom}_{gr}(Lx_i, Lx_i) \supseteq \prod_{\tau} K.$$

This leads to a contradiction since vLv and hence $\text{Hom}_{gr}(S, Lv)$ has K -dimension $\leq \sigma$ while, by Lemma 6.1, $\prod_{\tau} K$ has K -dimension $\geq 2^\tau > \sigma$. Hence Lv cannot be graded injective. ■

The proof of the next Lemma is essentially due Pere Ara. We gratefully thank him for offering such a short proof in place of our earlier elongated proof. In the proof we shall be using the following result which is well-known in the category of modules over an arbitrary ring R : If M is an injective left R -module, then $S/J(S)$ is self-injective, where S is the endomorphism ring of M and $J(S)$ is the Jacobson radical of S (see e.g. Theorem 13.1 in [23]). The same proof works for the graded version of this theorem.

Lemma 6.3 *If $v \in E^0$ and Lv is a graded injective left L -module, then v cannot be an infinite emitter.*

Proof. Suppose, by way of contradiction, the vertex v is an infinite emitter. Let $X = \{e_n : n = 1, 2, \dots\}$ be a countably infinite subset of edges emitted by v . Recall that the zero degree component L_0 of L is von Neumann regular (see Theorem 5.3, [10]). Consequently, vL_0v is von Neumann regular ring with identity and clearly its Jacobson radical $J(vL_0v) = 0$. Since Lv is graded injective, $vL_0v \cong \text{End}_L^{gr}(Lv)$ is a graded self-injective ring with identity (from the graded version of the proof of Theorem 13.1, [23]).

Let $S = \bigoplus_{i=1}^{\infty} vL_0ve_ie_i^* \subset vLv$. Let $\pi : S \rightarrow \bigoplus_{i=1}^{\infty} vL_0ve_{2i}e_{2i}^*$ be the coordinate

projection map so $\pi(e_{2i}e_{2i}^*) = e_{2i}e_{2i}^*$ for all i with $\text{Ker}(\pi) = \bigoplus_{i=1}^{\infty} vL_0ve_{2i+1}e_{2i+1}^*$.

Applying the Baer's criterion to the graded injective left module vL_0v , there is an element $a \in vL_0v$ (of degree 0) such that $\pi(x) = xa$. Let

$$a = \sum_{j=1}^m k_j p_j q_j^* \quad (\#)$$

where $k_j \in K$, the p_j, q_j are paths with $|p_j| = |q_j|$ and where we can assume that the monomials $p_j q_j^*$ are all different. First notice that it is not possible for $a = v$, since a satisfies the condition $e_1 e_1^* a = 0$. Also, we can assume, since $vav = a$, that $k_j v p_j q_j^* v = k_j p_j q_j^*$ for all $j = 1, \dots, m$. Let $T = \{e_n \in X : e_n^* p_j \neq 0 \text{ for some } p_j\}$. Clearly T is a finite set. We distinguish two cases.

Case 1: Suppose there is a term in $(\#)$ which is a multiple of v , say $k_i v$ with $k_i \neq 0$. Note that, by our assumption, no other term in $(\#)$ will then be a scalar multiple of v . As a consequence, for any $j \neq i$, the j -th term in $(\#)$ cannot be of the form $k_j q_j^*$ (without the real path p_j) or of the form $k_j p_j$ (without the ghost path q_j^*). Because, otherwise $\deg(q_j^*) = 0$ or $\deg(p_j) = 0$ so the j -th term will be of the form $k_j v$ for $j \neq i$, and this is impossible by our assumption.

Now for each odd integer $2n+1$, we have

$$0 = e_{2n+1} e_{2n+1}^* a = e_{2n+1} e_{2n+1}^* k_i v + e_{2n+1} e_{2n+1}^* \sum_{j=1, j \neq i}^m k_j p_j q_j^*.$$

If we choose $e_{2n+1} \notin T$, then $e_{2n+1}^* p_j = 0$ for all j and from the above equation we get, ■

$$-k_i e_{2n+1} e_{2n+1}^* = \sum_{j=1, j \neq i}^m k_j e_{2n+1} e_{2n+1}^* p_j q_j^* = 0, \text{ a contradiction.}$$

Case 2: Suppose, for each j , the j -th term in $(\#)$ contains the real path p_j .

Then for any $e_{2n} \notin T$, we have $e_{2n} e_{2n}^* = \pi(e_{2n} e_{2n}^*) = e_{2n} e_{2n}^* a = \sum_{j=1}^m k_j e_{2n} e_{2n}^* p_j q_j^* =$

0, again a contradiction. Hence v cannot be an infinite emitter.

Proposition 6.4 *If $L := L_K(E)$ is graded left self-injective, then (i) the graph E is row-finite and, (ii) for any vertex v , Lv cannot have infinitely many independent graded submodules.*

Proof. (i) If v is any vertex in E , then the direct summand Lv is an injective left L -module. By Lemma 6.3, v cannot be an infinite emitter. Hence the graph E must be row-finite. ■

(ii) To prove the second statement, let $v \in E^0$. Since E is row-finite, there are only finitely many paths of a fixed length n starting from v and so the set

Y of all finite paths α in E with $s(\alpha) = v$ is at most countable. Now every element of vLv is a finite sum of the form $\sum_{i=1}^n k_i p_i q_i^*$ where $k_i \in K$, p_i, q_i are finite paths with $s(p_i) = s(q_i) = v$ and $r(p_i) = r(q_i)$. Since the cardinality of the set of all finite subsets of the countable set Y is again countable, we conclude that the K -dimension of vLv is $\leq \aleph_0$. Since Lv is a graded injective L -module, we appeal to Lemma 6.2 to conclude that Lv cannot have infinitely many independent graded submodules.

Corollary 6.5 *If $L = L_K(E)$ is graded self-injective, then no cycle in E will have an exit*

Proof. Suppose, by way of contradiction, E contains a cycle c based at a vertex v and having an exit f at v . We modify the ideas in the proof of (ii) $\Rightarrow$ (iii) of Theorem 3.9 in [2]. Now for any n , $v - c^n(c^*)^n$ is a homogeneous idempotent and $v - c^n(c^*)^n = (v - c^n(c^*)^n)(v - c^{n+1}(c^*)^{n+1})$. Consequently we get an ascending chain of graded ideals contained in Lv :

$$L(v - cc^*) \subseteq L(v - c^2(c^*)^2) \subseteq \dots \quad (\#\#)$$

We claim that the containments in the chain $(\#\#)$ are strict. Suppose, on the contrary, $L(v - c^n(c^*)^n) = L(v - c^{n+1}(c^*)^{n+1})$ for some n . Then $(v - c^{n+1}(c^*)^{n+1}) = a(v - c^n(c^*)^n)$ for some $a \in L$. Multiplying the last equation by c^n on the right, we get $c^n - c^{n+1}c^* = 0$ and so $c^n = c^{n+1}c^*$. Again multiplying the last equation on the right by f , we get $c^n f = 0$, a contradiction. Thus $(\#\#)$ is a strictly increasing infinite chain of direct summands of Lv . Let, for each $n \geq 1$, $L(v - c^{n+1}(c^*)^{n+1}) = L(v - c^n(c^*)^n) \oplus B_n$. Then

$$A = \bigcup_{n \geq 1} L(v - c^n(c^*)^n) = L(v - cc^*) \oplus \bigoplus_{n=1}^{\infty} B_n \subset Lv.$$

Since Lv is graded-injective, this is a contradiction by Proposition 6.4. Thus no cycle in E has an exit. ■

Proposition 6.6 *If L is graded left self-injective, then every infinite path in E contains a line point or is tail-equivalent to a rational path induced by a cycle without exits and thus contains a Laurent vertex.*

Proof. Let μ be an infinite path in E with $s(\mu) = v$. Clearly Lv is a graded injective L -module. Suppose μ does not contain a line point. Then μ either contains a cycle or contains infinitely many bifurcating vertices. Suppose μ contains infinitely many bifurcating vertices v_n and bifurcating edges f_n with $s(f_n) = v_n$. Without loss of generality, we may assume $v = s(\mu) = v_1$. For each $n \geq 1$, let γ_n be the part of μ with $r(\gamma_n) = v_{n+1}$ and $s(\gamma_n) = v_n$ and thus μ is the concatenation of paths, $\mu = \gamma_1 \gamma_2 \cdots \gamma_n \cdots$. Let, for each n , $\epsilon_n = \gamma_1 \gamma_2 \cdots \gamma_n f_{n+1}^* f_{n+1}^* \gamma_n^* \cdots \gamma_1^*$. Then the set $\{\epsilon_n : n \geq 1\}$ is a set of homogeneous orthogonal idempotents giving rise to an infinite set of independent

graded submodules of Lv and this, by Proposition 6.4(ii), contradicts the graded injectivity of Lv . So μ must contain a cycle c which, by Corollary 6.5, has no exits. This means the infinite path μ must be tail-equivalent to the infinite rational path $ccc \cdots$ and contains the Laurent vertex v , where v is the base of the cycle c . ■

Theorem 6.7 *Let E be an arbitrary graph. Then the following are equivalent for $L := L_K(E)$:*

- (1) *L is graded left/right self-injective;*
- (2) *E is row-finite, no cycle in E has an exit and every infinite path in E is tail-equivalent to either straight line segment containing a line point or to a rational path $ccc \cdots$ for some cycle c (without exits);*
- (3) *There is a graded isomorphism*

$$L_K(E) \cong_{gr} \bigoplus_{v_i \in X} M_{\Lambda_i}(K)((|p^{v_i}|)) \oplus \bigoplus_{w_j \in Y} M_{\Upsilon_j}(K[x^{t_j}, x^{-t_j}])(|q^{w_j}|)$$

where Λ_i, Υ_j are suitable index sets, the t_j are positive integers, X is the set of representatives of distinct equivalence classes of line points in E and Y is the set of all distinct cycles (without exits) in E .

Proof. Now (1) $\Rightarrow$ (2) by Propositions 6.4, 6.6 and Corollary 6.5.

Assume (2). We claim that $E^0 = H$, the hereditary saturated closure of the set of all line points and Laurent vertices in E . Suppose, on the contrary, there is a $v_1 \in E^0 \setminus H$. Then, as E is row-finite, there must be an edge e_1 with $s(e_1) = v_1$ and $r(e_1) = v_2 \notin H$. This then implies there is an edge e_2 with $s(e_2) = v_2$ and $r(e_2) = v_3 \notin H$. Proceeding like this we get an infinite path $e_1 e_2 e_3 \cdots$ which neither contains a line point nor a cycle without exits (and thus no Laurent vertex), a contradiction to our hypothesis. Hence $E^0 = H$ and so L is the ideal generated by H . Then (3) follows from Theorem 2.10.

Finally, (3) $\Rightarrow$ (1) since L is graded semi-simple and so is graded injective as a left/right L -module. ■

References

- [1] G. Abrams, G. Aranda Pino, The Leavitt path algebra of a graph, J. Algebra 293 (2005) 319 - 334.
- [2] G. Abrams, G. Aranda Pino, F. Perera, M. Siles Molina, Chain conditions for Leavitt path algebras, Forum Math. 22 (2010) 95 - 114.
- [3] G. Abrams, G. Aranda Pino, M. Siles Molina, Finite dimensional Leavitt path algebras, J. Pure and Applied Algebra 209 (2007) 753 - 762.
- [4] G. Abrams, G. Aranda Pino, M. Siles Molina, Locally finite Leavitt path algebras, Israel J Math. 165 (2008) 329 - 348.

- [5] G. Abrams, J. Bell, K.M. Rangaswamy, On prime non-primitive von Neumann regular algebras, *Trans. Amer. Math. Soc.* 366 (2014) 2375 - 2392.
- [6] G. Abrams, F. Mantese, A. Tonolo, Extensions of simple modules over Leavitt path algebras, *J. Algebra* 431 (2015) 78 - 106.
- [7] G. Abrams, K.M. Rangaswamy, Regularity conditions for arbitrary Leavitt path algebras, *Algebr. Represent. Theory* 13 (2010) 319 - 334.
- [8] G. Abrams, K.M. Rangaswamy, M. Siles Molina, Socle series in a Leavitt path algebra, *Israel J Math* 184 (2011), 413 - 435.
- [9] P. Ara and M. Brustenga, Module theory over Leavitt path algebras and K-theory, *J. Pure and Applied Algebra* 214 (2010) 1131 - 1151.
- [10] P. Ara, M.A. Moreno, E. Pardo, Nonstable K-theory for Graph Algebras, *Algebr. Represent. Theory* 10 (2007) 157 - 168.
- [11] P. Ara and K.M. Rangaswamy, Finitely presented simple modules over Leavitt path algebras, *J. Algebra* 417 (2015) 333 - 352.
- [12] G. Aranda Pino, D. Martín Barquero, C. Martín Gonzalez and M. Siles Molina, The socle of a Leavitt path algebra *J. Pure and Applied Algebra* 212 (2008) 500 - 509.
- [13] G. Aranda Pino, K.M. Rangaswamy, M. Siles Molina, Weakly regular and self-injective Leavitt path algebras over arbitrary graphs, *Algebr. Represent. Theory* 14 (2011) 751 - 777.
- [14] X.W. Chen, Irreducible representations of Leavitt path algebras, *Forum Math.* 20 (2012), <http://dx.doi.org/10-1515/forum-2012-0020>.
- [15] A. Devulder, The speed of a branching system of random walks in random environment, *Statist. Probab. Lett.* 77 (18) (2007) 1712 - 1721.
- [16] D. Gonçalves, D. Royer, Unitary equivalence of representations of graph algebras and branching systems, *Functional Analysis and Applications* 45 (2011) 117 -127.
- [17] D. Gonçalves, D. Royer, Graph C^* -algebras, branching systems and the Perron-Frobenius operator, *J Math. Analysis and Applications* 391 (2012) 457 - 465.
- [18] D. Gonçalves, D. Royer, On the representations of Leavitt path algebras, *J. Algebra* 333 (2011) 258 - 272.
- [19] K.R. Goodearl, von Neumann regular rings, Krieger Publ. CO., Malabar (1991).
- [20] R. Hazrat, The graded structure of Leavitt path algebras, *Israel J. Math.* 195 (2013) 833 - 895.

- [21] R. Hazrat, Leavitt path algebras are graded von Neumann regular rings, *J Algebra* 401 (2014) 220 - 233.
- [22] K. J. Hochberg, Andreas Greven, On the use of the Laplace functional for two-level branching systems, *Int. J. Pure Appl. Math.* 55 (2) (2009) 165 - 172.
- [23] T.Y. Lam, *Lectures on Modules and Rings*, Graduate Texts in Math., vol. 189 Springer-Verlag, 1999.
- [24] C. Nastasescu, F. van Oystaeyen, *Graded Ring Theory*, North-Holland, Amsterdam, 1982.
- [25] K.M. Rangaswamy, Theory of prime ideals of Leavitt path algebras over arbitrary graphs, *J. Algebra* 375 (2013) 73 - 90.
- [26] K.M. Rangaswamy, On simple modules over Leavitt path algebras, *J. Algebra* 423 (2015) 239 - 258.
- [27] K.M. Rangaswamy, Leavitt path algebras with finitely presented irreducible representations, *arXiv:1501.01781 [math.RA]* 8 Jan 2015.
- [28] B. Salinier, R Strandh, Efficient simulation of forward-branching systems with constructor systems, *J. Symbolic Comput.* 22 (4) (1996) 381 - 399.
- [29] M. Tomforde, Uniqueness theorems and ideal structure for Leavitt path algebras, *J. Algebra* 318 (2007) 270 - 299.